



HUMAN GENOME EPIDEMIOLOGY (HuGE) REVIEW

Polymorphisms in Genes Involved in Folate Metabolism and Colorectal Neoplasia: A HuGE Review

Linda Sharp and Julian Little

From the Epidemiology Group, Department of Medicine and Therapeutics, University of Aberdeen, Aberdeen, Scotland.

Received for publication April 2, 2003; accepted for publication September 17, 2003.

Epidemiologic and mechanistic evidence suggests that folate is involved in colorectal neoplasia. Some polymorphic genes involved in folate metabolism—methylenetetrahydrofolate reductase (*MTHFR* C677T and A1298C), methionine synthase (*MTR* A2756G), methionine synthase reductase (*MTRR* A66G), cystathionine β -synthase (*CBS* exon 8, 68-base-pair insertion), and thymidylate synthase (*TS* enhancer region and 3' untranslated region)—have been investigated in colorectal neoplasia. For *MTHFR* C677T and A1298C, the variant allele is associated with reduced enzyme activity in vitro. For the other polymorphisms, functional data are limited and/or inconsistent. Genotype frequencies for all of the polymorphisms show marked ethnic and geographic variation. In most studies, *MTHFR* 677TT (10 studies, >4,000 cases) and 1298CC (four studies, >1,500 cases) are associated with moderately reduced colorectal cancer risk. In four of five genotype-diet interaction studies, 677TT subjects who had higher folate levels (or a "high-methyl diet") had the lowest cancer risk. In two studies, 677TT homozygote subjects with the highest alcohol intake had the highest cancer risk. Findings from six studies of *MTHFR* C677T and adenomatous polyps are inconsistent. There have been only one or two studies of the other polymorphisms; replication is needed. Overall, the roles of folate-pathway genes, folate, and related dietary factors in colorectal neoplasia are complex. Research priorities are suggested.

CBS; colorectal neoplasms; epidemiology; folic acid; *MTHFR*; *MTR*; *MTRR*; *TS*

Abbreviations: CBS, cystathionine β -synthase; CI, confidence interval; MSI, microsatellite instability; *MTHFR*, methylenetetrahydrofolate reductase; *MTR*, methionine synthase; *MTRR*, methionine synthase reductase; OR, odds ratio; rpt, repeat; *TS*, thymidylate synthase.

Editor's note: This article is also available on the Web site of the Human Genome Epidemiology Network (<http://www.cdc.gov/genomics/hugenet/default.htm>).

Evidence is accumulating for a role of folate in the etiology of colorectal carcinomas and adenomas (1). Many

of the genes involved in folate metabolism are polymorphic (2). This paper reviews five polymorphic genes—methylenetetrahydrofolate reductase (*MTHFR*), methionine synthase (*MTR*), methionine synthase reductase (*MTRR*), cystathionine β -synthase (*CBS*), and thymidylate synthase (*TS*)—and their associations with colorectal neoplasia.

Correspondence to Linda Sharp, Epidemiology Group, Department of Medicine and Therapeutics, University of Aberdeen, Polwarth Building, Foresterhill, Aberdeen AB25 2ZD, Scotland (e-mail: L.Sharp@abdn.ac.uk).




FIGURE 1. The roles of the methylenetetrahydrofolate reductase, methionine synthase, methionine synthase reductase, cystathionine β -synthase, and thymidylate synthase genes in the metabolism of folate.

GENES

5,10-MTHFR plays a central role in folate metabolism (figure 1), irreversibly converting 5,10-methylenetetrahydrofolate to 5-methylenetetrahydrofolate, the primary circulating form of folate. The substrate is vital for DNA synthesis. The product provides methyl groups for synthesis of methionine, a decreased pool of which may affect DNA methylation. The gene encoding 5,10-MTHFR, *MTHFR*, is located at 1p36.3 (3).

MTR, which is essential for maintaining adequate intracellular folate pools, catalyzes the remethylation of homocysteine to methionine, required for production of S-adenosylmethionine, the universal methyl group donor. Vitamin B₁₂ is a cofactor in this methylation process. The *MTR* gene is on 1q43 (4). MTR is maintained in its active form by MTRR (5), the gene for which, *MTRR*, is located at 5p15.3–p15.2. CBS catalyzes the conversion of homocysteine to cystathionine; vitamin B₆ is required in this reaction. The *CBS* gene is at 21q22.3. TS catalyzes the conversion of deoxyuridine monophosphate to thymidine monophosphate, requiring 5,10-methylenetetrahydrofolate as a methyl donor. The *TS* gene is located at 18p11.32.

Folate status could potentially be perturbed by polymorphisms in these genes. Two mechanisms have been proposed by which folate deficiency could affect malignancy: 1) by causing DNA hypomethylation and proto-oncogene activation and/or 2) by inducing uracil misincorporation during

DNA synthesis, leading to catastrophic DNA repair, DNA strand breakage, and chromosome damage (6). Human evidence in support of these mechanisms is limited (6, 7).

GENE VARIANTS

This section describes polymorphisms in the genes and their functional effects. With the exception of *MTHFR*, relatively few studies have investigated relations between the polymorphisms and blood levels of folate and related biomarkers in nondiseased persons. In subjects with medical conditions, it is possible that the condition or its treatment, rather than the underlying genotype, influences biomarker levels. Many studies have been small, with limited statistical power. A potential difficulty in interpretation is that any observed difference in biomarker levels by genotype may not be due to the polymorphism under study but to the presence of another polymorphism. Equally, a failure to observe differences in biomarkers by genotype could be due to the presence of another polymorphism with opposing functional effects. So far, there has been little investigation of the effects of combinations of polymorphisms. With regard to *MTHFR* C677T, only red cell folate measured by microbiologic assay is reliable; results of the radioimmune assay are biased (8). There is differential detection by the assays of various intracellular folates, the distribution of which is related to *MTHFR* genotype (9). Whether red cell folate

results measured by radioimmunoassay are biased for other polymorphisms in the folate-pathway genes is not known.

MTHFR

Several polymorphisms in the *MTHFR* gene have been reported, and two have been investigated in colorectal neoplasia: 1) C→T at nucleotide 677, leading to an alanine to valine conversion in the protein (10); and 2) A→C in exon 7, causing an alanine to glutamate protein change (11, 12). These polymorphisms are located 2.1 kb apart. The other polymorphisms—T1059C, T1317C, and G1793A (12–14)—are not discussed further in this paper.

For C677T, compared with homozygotes for the common variant (CC), heterozygotes have 65 percent of their enzyme activity levels in vitro and those who are homozygous variant (TT), 30 percent (15). From the microbiologic assay, compared with CC homozygotes, heterozygotes have 10 percent lower and TT homozygotes 18 percent lower red cell folate levels (16). Persons with the TT variant also have lowered plasma folate and vitamin B₁₂ levels and raised homocysteine levels (17, 18). In two studies, the association with homocysteine held only when folate status was low (19, 20); in another, it occurred only when riboflavin status was poor (21). Regarding *MTHFR* and DNA methylation, one small study found that DNA from subjects with the TT variant had a significantly higher methyl group acceptance capacity than DNA from subjects with the CC variant (22), but this finding was not confirmed in a larger study (23). In 292 subjects (66 percent of whom had coronary atherosclerosis) selected by *MTHFR* genotype (187 CC, 105 TT), DNA methylation status was affected by genotype among only those with lower plasma folate levels; subjects with the TT variant who had lower plasma folate concentrations had lower methylation levels than all other groups of subjects (24). A few studies have investigated *MTHFR* and uracil misincorporation, DNA strand breaks, or genetic instability in vivo and in vitro, with inconclusive results (23, 25–27).

For A1298C, enzyme activity in vitro is decreased in homozygotes variants (CC) and, to a lesser extent, in heterozygotes compared with those without the variant (11). Studies of A1298C and plasma folate and homocysteine are inconsistent (12, 28–31), which may be due to methodological reasons (e.g., non-population-based study, small sample size), or it may be that there is a relation that depends on the status of folate and/or related nutrients. Enzyme activity in vitro for compound heterozygotes (i.e., heterozygotes for C677T and for A1298C) is unclear (29).

MTR

The A-G polymorphism at position 2756 in the protein binding region of *MTR* replaces aspartic acid with glycine (32). Most studies suggest that plasma homocysteine level is lower in those with the rarer, G, than the more common, A, allele (18, 33–36). One study found significantly higher plasma folate levels in GG than in AA subjects (34), but this finding was not observed in another study (18). Evidence on red cell folate and on plasma vitamin B₁₂ and vitamin B₆ is very limited (18, 35, 37).

MTRR

The A66G polymorphism in the *MTRR* gene results in the substitution of isoleucine with methionine at codon 22 (5). In two studies, subjects homozygous for the common allele (AA) had elevated homocysteine levels compared with those who had other genotypes (38, 39); in a third study, genotype was not a significant predictor of homocysteine level (40). No associations were found between genotype and serum folate, vitamin B₆, or vitamin B₁₂ in the single known study (38).

CBS

Many mutations and several polymorphisms in the *CBS* gene have been reported (41). To our knowledge, the only variant investigated in colorectal neoplasia is the 68-base-pair insertion in the exon 8 coding region. Four studies found lower plasma homocysteine levels in persons carrying the insertion than in those without, although the difference was significant in only one (35, 36, 39, 42). One study suggested that the effect was modulated by plasma vitamin B₆ concentration (43); another suggested an interaction with *MTHFR* C677T (35). The one available study that we know of found no associations between genotype and red cell folate or plasma vitamin B₁₂ level (35).

TS

The *TS* enhancer region contains a series of 28-base-pair tandem repeats. Two repeats (2 rpt) or three repeats (3 rpt) are most common, with 3 rpt occurring most frequently. More repeats have been observed but are rare (44, 45). In vitro, compared with the double repeat, the triple repeat has been associated with 2.6-fold greater thymidylate synthase expression (46). Among 497 Singapore Chinese, plasma folate levels were significantly lower, and homocysteine levels nonsignificantly higher, in 3 rpt/3 rpt subjects than in those with other genotypes (47). When *MTHFR* and *TS* were considered together, plasma folate levels were highest (15.3 nM) in 677CC or 677CT and not 3 rpt/3 rpt subjects, intermediate (13.8 nM) in 677CC or 677CT and 3 rpt/3 rpt subjects, and lowest (11.6 nM) in 677TT subjects (irrespective of *TS* genotype).

The 3' untranslated region contains a 6-base-pair deletion at base pair 1494, the functional consequences of which are not known (48). The two polymorphisms appear to be in linkage disequilibrium (48).

Refer to the Appendix for Internet sites pertaining to the genes discussed in this review.

POPULATION FREQUENCIES

This section includes information on studies reporting genotype frequencies in persons without cancer or other diseases. Using appropriate Medical Subject Headings (MeSH) and text words, we searched MEDLINE, EMBASE, and PubMed databases for papers published from 1990 to December 2002. Further relevant articles were identified by hand-searching reference lists in published papers. *MTHFR*

frequencies are from the Human Genome Epidemiology (HuGE) reviews by Botto and Yang (49) and by Robien and Ulrich (50). The *A1298C* data reported by Robien and Ulrich are augmented with results from less-studied geographic areas and ethnic groups. For the studies tabulated here, Hardy-Weinberg equilibrium of the genotype frequencies was assessed by using the Pearson χ^2 test.

MTHFR

There is considerable ethnic and geographic variation in the frequency of the *C677T* variant (49). The *TT* prevalence ranged from around 1 percent in Black populations in the United States, sub-Saharan Africa, and South America to more than 20 percent in US Hispanics, Colombians, and Amerindians in Brazil. *TT* genotype frequency in White populations in Europe, North America, and Australia was 8–20 percent. In Europe, there appears to be a trend of increasing frequency of the variant from north to south. Twelve percent of Japanese were *TT* homozygotes.

For *A1298C*, the *CC* prevalence in North American studies, which included mainly White subjects, was 7–12 percent (50). In four Hispanic series ($n < 90$), the frequency was 4–5 percent (51–54). In two African-American series, 2 and 4 percent were *CC* subjects. In Europe, the prevalence of *CC* ranged from 4 to 12 percent in most studies. In two northeast Scotland series of subjects randomly selected from general practitioner registers, the frequencies were 15 percent (95 percent confidence interval (CI): 11.8, 19.2) and 18 percent (95 percent CI: 9.5, 30.4) (55, 56). In Chinese, Japanese, and Hawaiian populations, 1–4 percent were *CC* (50, 54) subjects. In the single studies in Brazil, Morocco, South Africa, and Turkey and among Israeli Jews, the frequencies were 6 percent (95 percent CI: 2.8, 9.6), 3 percent (95 percent CI not available), 4 percent (95 percent CI: 1.4, 9.9), 6 percent (95 percent CI: 1.7, 14.8), and 13 percent (95 percent CI: 9.7, 16.5), respectively (31, 33, 57–59).

In some series, but not all, a few persons with three or four variant alleles (i.e., *677TT/1298AC*, *CT/CC*, *TT/CC*) have been reported (35, 60–64).

MTR

In Japanese, Chinese, and Korean populations, the frequency of the *GG* genotype was 2–3 percent (18, 32–37, 54, 65–82; Web table 1). (This information is described in the first of four supplementary tables; each is referred to as “Web table” in the text and is posted on the Web site of the Human Genome Epidemiology Network (<http://www.cdc.gov/genomics/hugenet/default.htm>) as well as on the *Journal*'s Web site (<http://aje.oupjournals.org/>.) In most European series, approximately 3 percent of the subjects had the *GG* genotype. Frequencies from all but two North American studies were 1–5 percent. The frequency was 10–11 percent in these two series—one of White children and their mothers in Canada and the other of White persons in Hawaii. In the single African-American population, 6 percent (95 percent CI: 4.3, 8.7) of the subjects had the *GG* genotype. In

three studies, the genotype frequencies were not in Hardy-Weinberg equilibrium (73–75).

MTRR

The lowest reported prevalence of *GG* homozygotes was 8–10 percent in Japanese in Hawaii and in Hawaiians (5, 14, 38–40, 54, 83–85; Web table 2). Among 558 subjects in Northern Ireland, 12 percent (95 percent CI: 9.1, 14.6) were *GG* homozygotes, but this series was not in Hardy-Weinberg equilibrium. In most of the remaining series, the frequency was 19–29 percent. Among 97 African Americans and 96 Hispanics, the frequencies were 42 percent (95 percent CI: 32.3, 52.7) and 50 percent (95 percent CI: 39.6, 60.4), respectively.

CBS

Homozygosity for the 68-base-pair insertion is rare in all populations (35, 36, 39, 42, 54, 65, 70, 71, 77, 82, 86–98; Web table 3). The highest reported frequency was 3 percent among Blacks from Brazil and Africa. In four other series, the homozygote prevalence also reached 3 percent, but the genotype frequencies were not in Hardy-Weinberg equilibrium (42, 70, 71, 96). In Europe, Australia, and most US populations, the frequency of heterozygotes was 8–19 percent, with most around 13–15 percent. Two Japanese series found no heterozygotes. Heterozygosity occurred in 5 percent (95 percent CI: 1.6, 11.3) of the single Chinese series.

TS

In three studies in the United Kingdom, and in three of mainly White populations in the United States, 19–23 percent of subjects were 2 rpt/2 rpt (44–47, 99–102; Web table 4). The prevalence was 14–20 percent in two African and one African-American series and 17 percent among volunteers born in four southwest Asian countries living in Scotland. Two to 4 percent of two Chinese populations were homozygous variant. In all studies, genotype frequencies were in Hardy-Weinberg equilibrium.

In a single study of US Whites, 10 percent (95 percent CI: 7.7, 12.5) were homozygotes for the 3' untranslated region deletion (102).

Combinations of genotypes

Most studies reporting frequencies of combinations of genotypes are small (33, 35, 70, 80, 94, 103). In the largest, of almost 1,300 males in the United Kingdom, 8 percent carried the *CBS* 68-base-pair insertion and the *MTHFR* *T* allele; 5 percent of subjects had the *CBS* 68-base-pair insertion and the *MTR* *G* allele; and 20 percent carried both the *MTR* *G* and *MTHFR* *T* alleles (35).

Comments on studies of population frequencies

Few of the studies reviewed here were population based; many relied on convenience samples. Selection and partici-

pation biases may therefore explain some of the apparent variations in genotype prevalence. In a few studies, genotype frequencies were not in Hardy-Weinberg equilibrium. Although lack of Hardy-Weinberg equilibrium might indicate that the series were subject to selection or participation biases, there are other reasons why Hardy-Weinberg equilibrium might not hold, including migration or genotyping error (104). Many of the studies are relatively small, so the estimates of genotype frequency lack precision.

In many studies, the ethnic makeup of the participants is not described. Most well characterized are White populations in the United States and western Europe. Other populations, geographic areas, and ethnic groups, particularly in Africa, Asia (other than Japan), and South America, have been less studied. The generalizability from, for example, one "Black African" population to another may be limited since it is not always straightforward to establish ethnicity (105).

DISEASE

An estimated 945,000 new cases of colorectal cancer were diagnosed worldwide in 2000, and 492,000 persons died from the disease (106). Two thirds of incident cases occur in developed countries, where it is the third most common cancer in males and second most common in females (107). There are substantial international variations in incidence (108). Sixty to 70 percent of colorectal cancers arise in the colon (108).

Although most evidence is indirect, the majority of colorectal carcinomas are believed to develop from adenomatous polyps (109). Hyperplastic polyps may be precursors of some right-sided colon cancers (110). Investigation of the first occurrence, and the recurrence, of polyps may reveal factors important in early stages of the neoplastic process.

Fewer than 10 percent of incident colorectal tumors are due to hereditary nonpolyposis colorectal cancer and familial adenomatous polyposis (111). When these syndromes are excluded, there is still familial aggregation of cancers and adenomas (112–114), which is unlikely to be entirely accounted for by familial clustering of environmental factors (115). This information points to the potential importance of genetic susceptibility factors, and the interaction of these with each other and with environmental factors, in the disease causation.

The studies of Japanese migrants to the United States in the 1960s revealed the overwhelming importance of environmental factors in colorectal cancer etiology (116). Established risk factors for the disease are shown in table 1 (109, 117–125).

Although diet appears to be important in colorectal cancer (120), it has been difficult to identify the specific components involved. Observational epidemiologic evidence shows that a high vegetable intake is related to decreased risk (120), although recent work suggests that the relation is complex (124, 125). Vegetables, particularly green, leafy vegetables, are a major source of folate. The majority of prospective and case-control studies of serum folate, red cell folate, or reported dietary or total folate intake are compatible with inverse associations with colon cancer and

TABLE 1. Environmental factors associated with colorectal cancer

Increasing risk	Reducing risk
Excess weight*	Physical activity†
Tobacco smoking‡	Hormone replacement therapy§
Alcohol¶	Aspirin and other nonsteroidal antiinflammatory drugs#
	Vegetables**

* Bergström et al. (117); International Agency for Research on Cancer (IARC) Working Group (118).

† IARC Working Group (118).

‡ Giovannucci (119).

§ Beral et al. (121); Rossouw et al. (122).

¶ World Cancer Research Fund (WCRF)/American Institute for Cancer Research (AICR) (120); Cotton et al. (109): results of studies are heterogeneous.

IARC Working Group (123).

** WCRF/AICR (120); Terry et al. (124); Flood et al. (125): results of studies are heterogeneous.

adenomas (17, 54, 76, 125–146). There is no consistent association between rectal cancer and folate intake (126, 131, 133–135, 137, 138). One small trial of folic acid supplementation in persons from whom polyps had been removed observed a reduced recurrence rate in the supplemented group (147). Some studies are compatible with a positive association between alcohol intake, which adversely affects folate metabolism (148), and colorectal neoplasia (109). A "low-methyl diet," comprising high alcohol intake and low folate and methionine (and/or vitamins B₆ and B₁₂) intakes, has been associated with increased colon cancer risk (126, 130, 132, 140).

Internet sites providing data and information on colorectal neoplasia are contained in the Appendix.

ASSOCIATIONS

This section appraises studies of the polymorphisms and colorectal neoplasia risk. These studies were identified by using the search strategy described above with the addition of disease-specific Medical Subject Headings and text words.

MTHFR

C677T. To our knowledge, there have been 10 cancer studies: five in the United States, two in the United Kingdom, and one each in Australia, Mexico, and Korea (17, 54, 56, 98, 149–155; table 2). Two included only colon cancers (150, 154); the remainder included colon and rectal tumors. On the basis of the functional effects of the polymorphism, and the inverse association between folate status and disease, it might have been expected that the variant would be associated with increased disease risk. In contrast, seven studies were consistent with reduced risk in homozygous

TABLE 2. Studies of the *MTHFR* C677T genotype and colorectal carcinoma, with relative risks and 95% confidence intervals

Study area	Cases		Comparison group				Relative risk	95% CI*	Adjustment factors	Reference no.
	Type	No.	Type	No.	% TT	95% CI				
Australia†	Patients undergoing surgery for colorectal cancer at a hospital in Western Australia during 1985–1998; Duke's stage B or C; 46% male; 48% aged <70 years	501	"Healthy" persons from Western Australia; aged 20–92 years; 81% aged <70 years	1,207	11.0	8.4, 14.0	1.03‡	0.71, 1.49		98
Korea	Patients undergoing an operation for colorectal cancer at two centers; 51% male	200	"Healthy" unrelated adults without colorectal cancer; source not stated.	460	16.1	12.8, 19.8	0.75‡ 0.81‡	0.60, 0.95 0.46, 1.42		151
Mexico†	Patients with colorectal cancer	74	"Asymptomatic" subjects; source not stated	110	21.8	14.5, 30.7	0.94‡ 1.61‡	0.64, 1.39 0.62, 4.19		152
United Kingdom: Scotland	Residents of Grampian who had a first primary, histologically confirmed, colorectal cancer diagnosed in 1998–2000; 57% male; median age, 70 years	251	Persons randomly selected from lists of all those registered with general practitioners in Grampian; frequency matched to cases on age and sex; 51% male; median age, 62 years	394	11.9	8.9, 15.5	1.83‡ 0.93	0.84, 4.11 0.66, 1.32	Age, sex	56, 153
United Kingdom: Perth, Dundee, Leeds, York	Patients with incident colorectal cancer from four hospitals; aged 45–80 years; Caucasian; no history of familial adenomatous polyposis, inflammatory bowel disease, ulcerative colitis, diverticular disease, or previous malignancy	490	Controls from general practices; no history of previous cancer	592	8.3	6.2, 10.8	0.72 1.23	0.41, 1.28 0.81, 1.88		155
United States	Men enrolled in the Health Professionals Follow-up Study in 1986 who provided a blood sample in 1993–1994; self-reported colorectal cancer, confirmed from medical records and diagnosed in 1986–1994; aged 40–75 years at enrollment in 1986; cohort predominantly White	144	Male controls selected from the same cohort from among those who provided a blood sample in 1993–1994 but who did not report a diagnosis of colorectal cancer	627	13.4	10.8, 16.3	0.83 0.57	0.65, 1.07 0.30, 1.06	Age, family history, and intake of folate, methionine, and alcohol	149

United States	Male physicians participating in Physicians Health Study trial (exclusion criteria included history of myocardial infarction, stroke or ischemic heart disease, cancer, current renal or liver disease, peptic ulcer, or gout) who provided a blood sample at baseline in 1982 and reported colorectal cancer in 1982–1985, which was confirmed in medical records; mean age, 60 (standard deviation, 9) years	202	Male controls selected from the same cohort, matched to cases on age and smoking status; alive and free of colorectal cancer when matched case was diagnosed; mean age, 57 (standard deviation, 8) years	326	15.0	11.3, 19.4	TT vs. CC	0.45	0.24, 0.86	Age, smoking status, alcohol intake, multivitamin use, exercise, body mass index, aspirin use	17
United States: North Carolina	Persons with first invasive colon adenocarcinoma diagnosed in July 1996–June 2000, identified from cancer registry, aged 40–85 years at diagnosis, and had driver's license if under age 65 years; response rate, 66%; 52% male; 44% reported being African-American, 56% as White	552	Controls selected from 1) motor vehicle records (under age 65 years) or 2) lists of Medicare-eligible beneficiaries (aged ≥65 years); frequency matched to cases on ethnic group, age, sex; 38% African American, 62% White	868	6.6§	5.0, 8.4	CT vs. CC TT vs. CC	0.98 0.8	0.67, 1.45 0.5, 1.4	Age, ethnic group, sex, sampling fractions	154
United States: Utah and Minnesota	Participants in KPMCP* and residents of eight counties of Utah and Twin Cities area of Minnesota diagnosed with first primary colon cancer in 1991–1994; aged 30–74 years at diagnosis; 56% male; ethnic group of entire study population 4.2% Black, 4.4% Hispanic, 91.4% White; 75% of cases and controls genotyped	1,467	Controls 1) randomly selected from KPMCP lists, and 2) identified by random digit dialing and lists with driver's license or state identification in Minnesota and Utah (under age 65 years) and 3) randomly selected from Medical Care Financing lists in Utah (aged ≥65 years)	1,821	11.4	9.9, 12.9	CT vs. CC TT vs. CC	1.1 0.9	0.9, 1.4 0.7, 1.1	Age, body mass index, long-term vigorous physical activity, energy intake, dietary fiber, usual no. of cigarettes smoked	150
CT vs. CC									1.0	0.9, 1.2	Table continues

TABLE 2. Continued

Study area	Cases		Comparison group				Relative risk	95% CI	Adjustment factors	Reference no.
	Type	No.	Type	No.	% TT	95% CI				
United States: Hawaii	Persons with primary adenocarcinoma of the colon or rectum diagnosed in 1994–1998; identified through tumor registry; at least 75% Japanese or Caucasian or any percentage Hawaiian ancestry; 61% male; median age, 66 years; 59% Japanese, 27% Caucasian, 14% Hawaiian	548	Controls selected from 1) participants in an ongoing health survey among a 2% random sample of state households and 2) for those over age 65 years, Health Care Financing Administration rolls; 61% Japanese, 26% Caucasian, 13% Hawaiian	656	15.9 [¶]	13.1, 18.9	0.7	0.5, 1.0	Age, sex, ethnicity, smoking, physical activity, aspirin use, body mass index, schooling, intakes of nonstarch polysaccharides and calcium	54
							CT vs. CC	0.6, 1.1		

* MTHFR, methylenetetrahydrofolate reductase; CI, confidence interval; KPMCP, Kaiser Permanente Medical Care Program.

† DNA source: tumor for cases, blood for controls.

‡ Unmatched odds ratio, computed by Sharp and Little from data in the paper.

\$ % TT: African Americans = 1.8 (95% CI: 0.7, 3.9); Whites = 9.5 (95% CI: 7.1, 12.3).

¶ % TT: Japanese = 19.4 (95% CI: 15.6, 23.6); Caucasian = 14.0 (95% CI: 9.2, 20.2); Hawaiian = 3.4 (95% CI: 0.9, 9.6).

variant (*TT*) subjects compared with homozygotes for the common allele (17, 54, 149–151, 153, 154). Observed relative risks ranged from 0.45 to 0.9, although most did not reach statistical significance. A significant trend of decreasing risk with increasing number of *T* alleles has been reported (54). As has been observed in several meta-analyses of gene-disease associations (156, 157), the strongest effects were found in the two earliest studies (17, 149). Both were nested within cohort studies of predominantly White male populations in the United States. These populations were likely to have relatively high average intakes of total folate as a consequence of comparatively frequent use of vitamin supplements (158).

Although two studies were null overall (98, 155), one found an association with genotype in a subgroup (refer to the information later in this section; Shannon et al. (98)). In the other, although controls were matched to cases on age, sex, and general practice, this matching was not taken into account in the *MTHFR* analysis (155). The distribution by area of residence, which determines general practice, differed between cases and controls; if the prevalence of *MTHFR* variants differed between areas, this lack of adjustment could have affected the results. In addition, the *TT* prevalence among controls was lower than that in other studies from the United Kingdom.

In a study in Mexico, a nonsignificantly increased risk in carriers of the *T* allele was reported (152). This finding was based on small numbers of subjects, few details were provided about subject source populations, and the source of the DNA was tumor for cases and blood for controls.

One study observed that the inverse association with the *TT* genotype was stronger in older (aged 60–84 years) than in younger (aged 40–59 years) subjects, but this finding was not statistically significant (17). The same study reported that the inverse association held for tumors in both the colon and the rectum. In terms of location in the colon, Slaterry et al. (150) found that the *TT* genotype was associated with reduced risk in persons with proximal, but not those with distal, tumors. Two studies report results by ethnic group. Le Marchand et al. (54) found that the *TT* genotype was inversely associated with risk for subjects of Japanese origin and Caucasians, but not for Hawaiians. However, only nine Hawaiian subjects had the *TT* genotype. Keku et al. (154) found a modest inverse association among White subjects and African-American subjects.

Shannon et al. (98) stratified their cases into those showing microsatellite instability (MSI+) and those not (MSI–). *TT* genotype was associated with significantly raised risk in the MSI+ group (unadjusted odds ratio (OR) computed by us for *TT* vs. *CC* = 2.6, 95 percent CI: 1.08, 5.82) but not in the MSI– group. The MSI+ tumors were exclusively in the proximal colon and patients tended to be older, both factors that might have been expected to result in a reduced risk in *TT* subjects if the above observations regarding age and tumor location are true. This apparent inconsistency may be due to small numbers, bias, a failure to control for confounders, or chance. Further investigation to unravel the independent and joint influences of MSI, age, and tumor site is needed.

We know of six studies that have investigated *C677T* and adenomatous polyps, three in the United States and one each

in Japan, Norway, and Mexico (76, 152, 159–162; table 3). None found a significant association between genotype and risk, which raises the possibility that the *MTHFR* genotype may be relevant only in the later stages of the adenoma-carcinoma process, for example, in determining those persons with adenomas who will go on to develop carcinomas. It is also possible that the inconsistencies between the results of the studies of adenomas are due to differences between the studies in the subject source populations (i.e., whether they included screen-detected or symptomatic adenomas) and in the control series (e.g., whether it comprised polyp-free subjects).

In two studies of hyperplastic polyps, no association was found between genotype and disease (162, 163; table 4).

A1298C. Four studies, three in the United States (28, 54, 154) and one in Scotland (56), have investigated the role of *A1298C* in cancer (table 5). In all, risk was modestly reduced in *CC* compared with *AA* subjects. Relative risks were in the range of 0.6–0.8 and mostly did not reach statistical significance. Since this finding is consistent with the pattern observed for *C677T*, it raises the possibility that the *A1298C*-cancer relation is actually due to *C677T*. However, Chen et al. (28) reported that the *A1298C* result was not due to confounding by *C677T*. In addition, Le Marchand et al. (54) found that, compared with *677CC/1298AA* persons, those who carried *677T* and *1298C* had the lowest risk. Keku et al. (154) reported that the *A1298C*-cancer association was stronger among White than African-American subjects.

MTR

One cancer study and one of adenomas found a slightly reduced risk for *GG* homozygotes (18, 76; table 5). A third study found no effect overall but observed an inverse association between *GG* and cancer among a subgroup of Hawaiian subjects (54).

MTRR

In the single study that we know of, in Hawaii, *A66G* was not associated with cancer when the three ethnic groups included in the study were analyzed together (54; table 5). However, among White subjects, there was a trend of borderline significance of increasing risk with increasing number of variant alleles (OR for *GG* vs. *AA* = 1.9, 95 percent CI: 1.0, 3.8; *p* for trend = 0.07).

CBS

Heterozygotes for the *CBS* insertion were twice as frequent among controls as among cancer cases in one study (OR computed by us = 0.50, 95 percent CI: 0.24, 1.07) (98; table 5). Compatible with this finding, the other available study suggested that the variant was associated with reduced cancer risk (54).

TS

In the single study that we are aware of, of the 6-base-pair deletion and cancer in non-Hispanic White subjects in the United States, which was reported in abstract form only, subjects with the deletion had a relative risk of 1.40 (95 percent CI: 0.99, 1.98; *p* = 0.058) compared with those with no deletion allele (164; table 5). In another study of men in the United States, again reported only as an abstract, 2 rpt homozygous persons had a nonsignificantly reduced cancer risk (relative risk for 2 rpt/2 rpt vs. 3 rpt/3 rpt = 0.65, 95 percent CI: 0.38, 1.12) (99). In the single study of adenomas, no significant association was found between either polymorphism and disease, nor did combinations of the two polymorphisms affect risk (102).

Other diseases

Genetic variation in *MTHFR*, *CBS*, *MTR*, *MTRR*, and *TS* has been investigated in other conditions in which folate or homocysteine may be involved. Examples are congenital anomalies such as neural tube defects, Down's syndrome, and orofacial clefts (5, 40, 49, 84, 165, 166); cancers including leukemia and lymphomas, breast, gastric, and esophageal tumors (50, 55, 64, 67, 167); cardiovascular disease (34, 87, 158, 168, 169); and Alzheimer's disease (170).

INTERACTIONS

Gene-environment interactions

MTHFR C677T. The gene-environment interactions explored have concerned features of the "low-methyl" diet and genotype. Four of five studies suggest interactions between folate, methionine, or alcohol and *C677T* in relation to cancer. Chen et al. (149) reported that the inverse association with the *TT* genotype was greatest among persons in the highest tertiles of folate and methionine intake. The results of Ma et al. (17), who examined plasma folate, and Le Marchand et al. (54), who analyzed food and total folate intake, were compatible with this finding. Keku et al. (154), however, did not observe this pattern with regard to total folate intake.

Slattery et al. (150) categorized subjects as consuming low-, intermediate-, and high-methyl diets. The lowest odds ratio was for subjects with the *TT* genotype consuming a high-methyl diet (OR for high-methyl and *TT* vs. low-methyl and *CC* = 0.4, 95 percent CI: 0.1, 0.9), while the odds ratios for subjects consuming a low-methyl diet did not vary by genotype (150). Consistent with this finding, Ma et al. (17) observed an increased risk among the folate deficient (plasma folate <3.0 ng/ml) irrespective of genotype.

Two cancer studies found significant interactions between *C677T* and alcohol (17, 149). High intake abolished the reduced risk associated with the *TT* genotype to the extent that subjects with this *TT* genotype who consumed the largest quantities of alcohol were at the greatest risk of cancer (greater even than for those without the *T* allele who

TABLE 3. Studies of the *MTFRR** *C677T* genotype and adenomatous polyps, with relative risks and 95% confidence intervals

Study area	Cases		Comparison group				Comparison	Relative risk	95% CI*	Adjustment factors	Reference no.
	Type	No.	Type	No.	% TT	95% CI					
United States	Women enrolled in the Nurses' Health Study in 1976 who provided a blood sample in 1989–1990; first incident proximal or distal colorectal adenoma diagnosed during time from blood specimen to June 1994; approximately 95% of the cohort is White	257	1) Cohort members in whom colorectal adenoma had not been diagnosed and who were born in the same year as the matched case and had had a sigmoidoscopy since the blood sample was taken ($n = 257$); plus 2) female cohort members who had served as controls for a breast cancer study, of whom 71% had not had a sigmoidoscopy and 29% had had a sigmoidoscopy and did not have an adenoma	713	9.3	7.2, 11.6	TT vs. CT/CC	1.35	0.84, 21.7	Age, family history, smoking status, body mass index, and intakes of folate, methionine, alcohol, fiber, and saturated fat	76
United States: California	Subjects undergoing screening sigmoidoscopy at two medical centers during 1991–1993; aged 50–74 years; no evidence of prior bowel disease and no previous bowel surgery	471	Without any adenoma at sigmoidoscopy and no history of adenomas; matched to cases on sex, sex, date of sigmoidoscopy, and clinic	510	9.6	7.2, 12.5	TT vs. CC	1.11	0.71, 1.71	Age, race, sex, clinic, date of sigmoidoscopy	160
United States: Minneapolis, Minnesota	Subjects recruited from private gastroenterology practice undertaking colonoscopies in 10 hospitals; underwent colonoscopy in 1991–1994; English speaking; no known genetic syndromes predisposing to colorectal cancer; no history of cancer or inflammatory bowel disease; aged 30–74 years; participation rate, 68%	527	Free of all polyps at colonoscopy; 38% male; mean age, 52.8 (standard deviation, 10.9) years; 97% White	645	11.0	8.7, 13.7	TT vs. CC	0.85	0.65, 1.13	Age, sex, body mass index, use of hormone replacement therapy, and percentage of calories from fat, dietary fiber, folate, vitamin B ₁₂ , vitamin B ₆ , methionine, alcohol	159
Japan	Male military officials undergoing preretirement health examination at two hospitals; had a partial or total colonoscopy and provided a blood sample; aged 47–55 years; no prior history of colectomy, polypectomy, malignant neoplasia	205	Normal total colonoscopy without in situ or invasive carcinoma	220	11.8	7.9, 16.8	TT vs. CC	0.87	0.56, 1.34	Hospital, employment, military rank, smoking, alcohol intake	161
							CT vs. CC	1.17	0.61, 2.23		

Norway	Participants in Telemark I study; born in 1924–1933; selected from population register in 1983 and randomly assigned to endoscopy or control group; 799 participated; in 1996, offered colonoscopy and removal of polyps; results available for 443 participants (229 male, 214 female; median age, 67 years)	47	Without polyps ($n = 116$) or with hyperplastic polyps or "low-risk" adenomas ($n = 278$)	394	7.1	4.8, 10.1	TT vs. CC	2.41	0.82, 7.06	Age, sex, red blood cell folate, use of nonsteroidal antiinflammatory drugs, flexible sigmoidoscopy in 1983, body mass index, current smoking	162
Mexico	Subjects with "high-risk" colorectal adenomas (≥ 10 mm or severe dysplasia or villous components)	32	"Asymptomatic" subjects; source not stated	110	21.8	14.5, 30.7	CT vs. CC	1.51	0.76, 2.99		
	Patients with colorectal adenomas						TT vs. CC	1.65†	0.41, 6.73		152
							CT vs. CC	0.98†	0.28, 3.67		

* MTHFR, methylenetetrahydrofolate reductase; CI, confidence interval.

† Unmatched odds ratio, computed by Sharp and Little from data in the paper.

were in the highest alcohol group). Keku et al. (154) found no interaction with alcohol but did not consider quantity, only whether subjects had "ever" or "never" consumed alcohol.

High blood riboflavin levels may improve MTHFR activity in *TT* persons because the cofactor for MTHFR is a metabolite of riboflavin (171). Le Marchand et al. (54) observed the lowest relative risk for cancer among *TT* persons with the highest riboflavin intake. Genotype-folate-riboflavin combinations were not considered.

Little is published on gene-diet interactions and adenomas. In the two known studies, the stratum of highest risk comprised *TT* persons with the lowest red cell or plasma folate levels (160) or the lowest intakes of folate, methionine, vitamin B₆, and vitamin B₁₂ (159), but the gene-nutrient interactions were not statistically significant. With regard to alcohol and genotype, the pattern observed is similar to that for cancer (159, 160).

MTHFR A1298C. Keku et al. (154) observed a significant interaction ($p = 0.03$) between total folate intake and *A1298C* genotype among White but not African-American subjects; fewer African-American subjects were involved in the study. Unlike the pattern for *C677T*, White *1298CC* subjects who consumed less than 400 ng of folate per day had a greater reduced cancer risk than those whose folate intake was higher. No interactions were observed between *A1298C* and "ever" or "never" consuming alcohol.

Two further studies of *A1298C* reported no significant interactions with blood levels or intake of folate or related nutrients and colorectal neoplasia (28, 54). The results were not shown.

MTR. For cancer, Ma et al. (18) reported a significant interaction between *MTR* and alcohol intake (table 5); persons with the *GG* genotype consuming more than one drink a day had an increased disease risk (OR for *GG* and ≥ 1 drink/day vs. *AA* and < 1 drink/day = 2.64, 95 percent CI: 0.65, 10.82), while those consuming less than one drink a day had a reduced risk (OR = 0.27, 95 percent CI: 0.09, 0.81; p for interaction = 0.04). There was also a nonsignificant 50 percent risk reduction among *GG* subjects whose plasma folate levels were in the upper two tertiles compared with those with the same folate level and the *AA/AG* genotype; persons with the *GG* genotype in the lowest plasma folate tertile did not have a reduced risk (p for interaction = 0.22).

MTRR and CBS. Le Marchand et al. (54) reported no significant interactions between *MTRR* or *CBS* and dietary folate, vitamin B₁₂, vitamin B₆, riboflavin, or methionine. Results were not shown.

TS. For adenomas, Ulrich et al. (102) found a statistically significant interaction between the tandem repeat polymorphism and folate intake. Among 3 rpt/3 rpt persons, higher folate intake (> 440 ng/day) was associated with a 50 percent reduced risk compared with lower folate intake. However, among 2 rpt/2 rpt persons, higher folate intake was associated with a 50 percent increased risk (p for interaction = 0.03). A similar pattern was observed for vitamin B₁₂ intake (p for interaction = 0.08). No interactions were found with intakes of vitamin B₆, methionine, or alcohol, nor were there interactions between the 3' untranslated region polymorphism and dietary variables.

TABLE 4. Studies of the *MTHFR** *C677T* genotype and hyperplastic polyps, with relative risks and 95% confidence intervals

Study area	Cases		Comparison group				Comparison	Relative risk	95% CI*	Adjustment factors	Reference no.
	Type	No.	Type	No.	% <i>TT</i>	95% CI					
Norway	Participants in Telemark I study; born 1924–1933; selected from population register in 1983 and randomly assigned to endoscopy or control group; 799 participated; in 1996, offered colonoscopy and removal of polyps; results available for 443 participants (229 male, 214 female; median age, 67 years)										162
	With "high-risk" hyperplastic polyps (<i>n</i> ≥ 3)	91	Without polyps (<i>n</i> = 116) or with adenomas or "low-risk" hyperplastic polyps (<i>n</i> = 233)	349	7.1	4.8, 10.1	<i>TT/CT</i> vs. <i>CC</i>	1.43†	0.87, 2.33		
United States: Minneapolis, Minnesota	Subjects recruited from private gastroenterology practice undertaking colonoscopies in 10 hospitals; underwent colonoscopy in 1991–1994; English speaking; without known genetic syndromes predisposing to colorectal cancer; no history of cancer or inflammatory bowel disease; aged 30–74 years										163
	Diagnosis of colon or rectal hyperplastic polyps; 97% White; 57% male; mean age, 53.7 years	200	Free of all polyps at colonoscopy; 97% White; 38% male; mean age, 52.8 (standard deviation, 10.9) years	645	11.0	8.7, 13.7	<i>TT</i> vs. <i>CC</i>	0.9	0.5, 1.6	Age, sex, body mass index, use of hormone replacement therapy, smoking, percentage of calories from fat, dietary fiber, folate, vitamin B ₁₂ , vitamin B ₆ , methionine, alcohol	
							<i>CT</i> vs. <i>CC</i>	0.8	0.6, 1.2		

* *MTHFR*, methylenetetrahydrofolate reductase; CI, confidence interval.

† Unmatched odds ratio, computed by Sharp and Little from data in the paper.

Gene-gene interactions

Metabolism of any exposure is likely to depend on the balance between the relative activities of all of the enzymes active within the metabolic pathway (172). So far, we know of two studies that have considered joint effects of folate-pathway genes (54, 102; table 5).

For cancer, Le Marchand et al. (54) observed that the *MTHFR* *T* allele had the greatest effect among subjects with the *MTR* *G* allele (OR for *CT/TT* and *AG/GG* vs. *CC* and *AA* = 0.7, 95 percent CI: 0.5, 1.0; p for interaction = 0.05). Considering *MTHFR* *C677T* and *CBS*, they reported that the group with both variants appeared to be at reduced risk; however, this result was based on small numbers, and the interaction was not significant. Meanwhile, *MTRR* did not interact with *MTHFR* *C677T*.

For adenomas, Ulrich et al. (102) investigated interactions between *C677T*, *TS* tandem repeat, and folate intake. The association of higher folate intake with reduced risk among 3 rpt/3 rpt subjects was not modified by *MTHFR*. The increased risk associated with lower folate intake in *TT* subjects appeared limited to 3 rpt homozygotes. These findings were not statistically significant.

Comments on studies of gene-disease associations and interactions

Some of the heterogeneity in the findings with regard to the genotype main effects is likely to be due to differences between the populations studied in average levels of intake of folate, alcohol, and related dietary factors. If there truly are interactions between genotype and folate, for example, they may be seen only in populations with high or low folate levels (depending on the direction of the interaction). Such an effect has recently been observed for *MTHFR* *C677T* and coronary heart disease (158).

Methodological factors are also important. Five cancer studies (17, 56, 149, 151, 152) and four adenoma studies (76, 152, 161, 162) each included fewer than 300 cases and thus had limited statistical power, particularly for subgroup and interaction analyses. The nonprospective studies are most susceptible to bias. Some were not population based. In some, it is not clear whether the controls came from the population that gave rise to the cases. In others, the case series were limited to subjects still alive to provide a DNA sample (prevalent cases), which would have resulted in bias if any of the genotypes were associated with survival (currently not known). Few studies provided information on

participation rates, making it difficult to assess bias and generalizability. It is likely that a proportion of the controls in the cancer studies may have been harboring undiagnosed polyps. Depending on the relations between each polymorphism and adenomas, this may have introduced random error or bias. The presence of undetected polyps among controls would not be important if the genotype was etiologically relevant only *after* an adenoma had developed, as seems likely for *MTHFR C677T*. For the other genotypes, it is not clear at what stage in the adenoma-carcinoma sequence they may be relevant. Finally, the possibility cannot be discounted that the findings do not reflect an association between the specified polymorphisms and colorectal neoplasia but rather are a consequence of linkage disequilibrium.

LABORATORY TESTS

MTHFR C677T and *A1298C* are detected by means of DNA amplification using polymerase chain reaction followed by restriction fragment length polymorphism analysis; *HinfI* for *C677T* and *MboII* (12) for *A1298C* (10, 11) are used. The *MTR* and *MTRR* polymorphisms and the 3' untranslated region variant in *TS* are also detected by restriction fragment length polymorphism, with digestion with *MaeII* for *MTR*, with *NdeI* or *AflIII* for *MTRR*, and with *DraI* for *TS* (4, 5, 48, 54). The *TS* tandem repeat and *CBS* insertion are detected by DNA amplification and visualization on agarose gels (46, 97).

Most studies did not report the success rate in extracting DNA from samples, the proportion of eligible subjects for whom genotyping failed, or the degree of genotyping reproducibility, all of which are important indicators of the analytical validity of genotyping (173).

Laboratories are increasingly using high-throughput genotyping methods, an area of considerable development and innovation. Although quality control and analytical validity in this context are important (173), published data are currently lacking.

POPULATION TESTING

Companies in the United States and the United Kingdom are offering consumer tests for genotypic or phenotypic markers of polymorphisms influencing nutrient metabolism, including *MTHFR* (174, 175). However, the scientific evidence currently is not strong enough to advocate population testing for any polymorphisms reviewed here.

Testing for these polymorphisms might be valuable in cancer patients. 5-Fluorouracil, commonly used in colorectal cancer chemotherapy, is a thymidylate synthase inhibitor and can cause severe folate depletion. Knowledge of patient genotype could be used to tailor chemotherapy regimes to 1) minimize toxicity and side effects, thus improving quality of life, and/or 2) increase the effectiveness of treatment and ultimately lengthen survival. So far, evidence in this area is limited to the *TS* tandem repeat and *MTHFR C677T*. Among 51 stage III colon cancer patients treated with 5-fluorouracil and leucovorin (folinic acid), presence of the *MTHFR T*

allele had little effect on probability of death or length of survival in those who had died, except in 12 patients with rectosigmoid colon cancer (176). In a study of 365 nonadjuvant-treated patients, the *TT* genotype was associated with improved survival, but this result did not persist after adjustment for disease stage (98).

For *TS*, some (177–179) but not all (180, 181) studies of colorectal cancer patients concluded that higher *TS* tumor expression levels were related to shorter survival. Consistent with this finding, one genotype study suggested that carrying the 3 rpt allele increased risk of death (179). Four studies of genotype and response to 5-fluorouracil (182–185) suggested that 2 rpt/2 rpt patients may be more responsive to therapy but subject to greater toxicity (186). Most of the studies (of genotype or phenotype) have been small, included selected patient groups, and made limited adjustment for potentially important factors.

CONCLUSIONS AND RESEARCH PRIORITIES

The observed association of the *MTHFR* homozygous variant genotypes with reduced carcinoma risk was the opposite of what might have been expected a priori. This finding has led investigators to reconsider the folate metabolism pathway, putting a greater emphasis on the functions of folate and *MTHFR* in DNA synthesis. The evidence is compatible with interactions between *MTHFR* genotype and folate, alcohol, and/or related nutrients in relation to colorectal cancer. Evidence on polymorphisms other than *MTHFR C677T* is extremely limited. The associations observed between *MTR*, *CBS*, *MTRR*, and *TS* genotypes and colorectal neoplasia are tentative at best and require replication. The few studies of combinations of polymorphisms suggest the possibility of gene-gene interactions; again, further investigation is needed to confirm initial findings. Altogether, the evidence suggests that the roles of folate-metabolizing genes, folate, and related dietary factors in colorectal neoplasia are complex. Methodologies are currently lacking for specification of hypotheses, clarification of functional effects, and statistical analysis relating to such complex gene-environment pathways. This area of research must be a priority if advancements in understanding of disease etiology are to be achieved. Table 6 lists other areas for further research.

REFERENCES

1. Little J, Sharp L. Colorectal neoplasia and genetic polymorphisms associated with folate metabolism. *Eur J Cancer Prev* 2002;11:105–10.
2. Johnson WG. DNA polymorphism-diet-cofactor-development hypothesis and the gene-teratogen model for schizophrenia and other developmental disorders. *Am J Med Genet* 1999;88:311–23.
3. Goyette P, Sumner JS, Milos R, et al. Human methylenetetrahydrofolate reductase: isolation of cDNA, mapping and mutation identification. *Nat Genet* 1994;7:195–200.
4. Leclerc D, Campeau E, Goyette P, et al. Human methionine synthase: cDNA cloning and identification of mutations in

TABLE 5. Summary of studies of the *MTHFR** *A1298C* polymorphism, other folate pathway genes, and colorectal neoplasia

Gene	Polymorphism	Study area, study design, cases†	Gene-disease associations			Gene-environment interaction	Gene-gene interactions	Reference no.
			Comparison	Relative risk	95% CI*			
<i>MTHFR</i>	<i>A1298C</i>	United States, case-control, carcinoma	CC vs. AA	0.8	0.5, 1.4	No interactions with total or dietary folate, vitamin B ₆ , vitamin B ₁₂ , riboflavin, methionine, ethanol‡		54
		United States, nested case-control, carcinoma	CC vs. AA	0.73	0.37, 1.43	Risk associated with CC not modified by plasma folate status‡		17, 28
		United States, case-control, carcinoma	CC vs. AA	0.6	0.4, 0.9	Significant interaction between <i>A1298C</i> and total folate intake for Whites only; among African Americans, combined <i>C677T</i> and <i>A1298C</i> genotype and total folate produced interaction of borderline significance; no significant interactions of <i>A1298C</i> and alcohol intake for either ethnic group		154
		Scotland, case-control, carcinoma	CC vs. AA	0.67	0.39, 1.13	—§		56
<i>MTR</i> *	<i>A2756G</i>	United States, nested case-control, adenoma	GG vs. AA	0.66	0.26, 1.70	—§	No significant interaction with <i>MTHFR C677T</i> ‡	76
		United States, case-control, carcinoma	GG vs. AA	1.1	0.6, 2.2	No interactions with total or dietary folate, vitamin B ₆ , vitamin B ₁₂ , riboflavin, methionine, ethanol‡	Significant interaction with <i>MTHFR C677T</i>	54
		United States, nested case-control, carcinoma	GG vs. AA	0.59	0.27, 1.27	Significant interaction with alcohol; suggestion of possible joint effect with plasma folate, but not significant; no interaction with homocysteine; no significant interaction with vitamin B ₁₂ ‡	—§	18

Table continues

- patients of the cblG complementation group of folate/cobalamin disorders. *Hum Mol Genet* 1996;5:1867–74.
- Wilson A, Platt R, Wu Q, et al. A common variant in methionine synthase reductase combined with low cobalamin (vitamin B₁₂) increases risk for spina bifida. *Mol Genet Metab* 1999;67:317–23.
 - Duthie SJ. Folic acid deficiency and cancer: mechanisms of DNA instability. *Br Med Bull* 1999;55:578–92.
 - Ames BN. DNA damage from micronutrient deficiencies is likely to be a major cause of cancer. *Mutat Res* 2001;475:7–20.
 - Molloy AM, Mills JL, Kirke PN, et al. Whole-blood folate values in subjects with different methylenetetrahydrofolate reductase genotypes: differences between the radioassay and microbiological assays. *Clin Chem* 1998;44:186–8.
 - Bagley PJ, Selhub J. A common mutation in the methylenetetrahydrofolate reductase gene is associated with an accumulation of formylated tetrahydrofolates in red blood cells. *Proc Natl Acad Sci U S A* 1998;95:13217–20.
 - Frosst P, Blom HJ, Milos R, et al. A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. *Nat Genet* 1995;10:111–13.
 - van der Put NMJ, Gabreëls F, Stevens EMB, et al. A second common mutation in the methylenetetrahydrofolate reductase gene: an additional risk factor for neural-tube defects? *Am J Hum Genet* 1998;62:1044–51.
 - Weisberg I, Tran P, Christensen B, et al. A second genetic polymorphism in methylenetetrahydrofolate reductase (*MTHFR*) associated with decreased enzyme activity. *Mol Genet Metab* 1998;64:169–72.
 - Trembath D, Sherbondy AL, Vandyke DC, et al. Analysis of select folate pathway genes, *PAX3*, and human *T* in a midwestern neural tube defect population. *Teratology* 1999;59:331–41.
 - Rady PL, Szucs S, Grady J, et al. Genetic polymorphisms of methylenetetrahydrofolate reductase (*MTHFR*) and methionine synthase reductase (*MTRR*) in ethnic populations in Texas; a report of a novel *MTHFR* polymorphic site, G1793A. *Am J Med Genet* 2002;107:162–8.
 - Rozen R. Genetic predisposition to hyperhomocysteinemia: deficiency of methylenetetrahydrofolate reductase (*MTHFR*). *Thromb Haemost* 1997;78:523–6.
 - Molloy AM, Daly S, Mills JL, et al. Thermolabile variant of 5,10-methylenetetrahydrofolate reductase associated with low

TABLE 5. Continued

Gene	Polymorphism	Study area, study design, cases†	Gene-disease associations			Gene-environment interaction	Gene-gene interactions	Reference no.
			Comparison	Relative risk	95% CI*			
<i>MTRR</i> *	<i>A66G</i>	United States, case-control, carcinoma	<i>GG</i> vs. <i>AA</i>	1.4	0.9, 2.0	No interactions with total or dietary folate, vitamin B ₆ , vitamin B ₁₂ , riboflavin, methionine, ethanol‡	No interaction with <i>MTHFR C677T</i>	54
<i>CBS</i> *	68 bp* insertion	United States, case-control, carcinoma	Weak inverse association with presence of insertion¶			No interactions with total or dietary folate, vitamin B ₆ , vitamin B ₁₂ , riboflavin, methionine, ethanol‡	Weak suggestion of possible interaction with <i>MTHFR C677T</i>	54
		Australia, case-control, carcinoma#	Frequency of heterozygotes in controls (10%) vs. cases (5%)			—§		98
<i>TS</i> *	28 bp tandem repeat	United States, case-control, adenoma	2 rpt/2 rpt vs. 3 rpt/3 rpt	0.9	0.6, 1.3	Significant interaction with total folate intake; borderline significant interaction with total vitamin B ₁₂ intake	Suggestion of joint effect with <i>MTHFR C677T</i>	102
	28 bp tandem repeat	United States, case-control, carcinoma	2 rpt/2 rpt vs. 3 rpt/3 rpt	0.65	0.38, 1.12	No interactions with vitamin B ₆ , methionine, or alcohol‡		99
	6 bp deletion in 3' untranslated region	United States, case-control, carcinoma	With deletion vs. no deletion	1.40	0.99, 1.98			164
	6 bp deletion in 3' untranslated region	United States, case-control, adenoma	Homozygous no deletion vs. 6 bp/6 bp	1.13	0.73, 1.74	No consistent patterns with dietary folate or vitamin B ₁₂ ‡	No consistent patterns with <i>MTHFR C677T</i> ‡	102

* *MTHFR*, methylenetetrahydrofolate reductase; CI, confidence interval; *MTR*, methionine synthase; *MTRR*, methionine synthase reductase; *CBS*, cystathionine β-synthase; bp, base pair; *TS*, thymidylate synthase; rpt, repeat.

† Refer to tables 2 and 3 for further details of the study populations, etc.

‡ Data not shown in the paper.

§ None mentioned in the paper as having been investigated.

¶ Unadjusted odds ratio for heterozygotes or homozygous insertion vs. homozygous no insertion = 0.88 (95% CI: 0.50, 1.55).

This analysis included only 155 of the original control series.

red-cell folates: implications for folate intake recommendations. *Lancet* 1997;349:1591–3.

17. Ma J, Stampfer MJ, Giovannucci E, et al. Methylenetetrahydrofolate reductase polymorphism, dietary interactions and risk of colorectal cancer. *Cancer Res* 1997;57:1098–102.
18. Ma J, Stampfer MJ, Christensen B, et al. A polymorphism of the methionine synthase gene: association with plasma folate, vitamin B₁₂, homocyst(e)ine, and colorectal cancer risk. *Cancer Epidemiol Biomarkers Prev* 1999;8:825–9.
19. Jacques PF, Bostom AG, Williams RR, et al. Relation between folate status, a common mutation in methylenetetrahydrofolate reductase, and plasma homocysteine concentrations. *Circulation* 1996;93:7–9.
20. Girelli D, Friso S, Trabetti E, et al. Methylenetetrahydrofolate reductase C677T mutation, plasma homocysteine, and folate in subjects from northern Italy with or without angiographically documented severe coronary atherosclerotic disease: evidence for an important genetic-environmental interaction. *Blood* 1998;91:4158–63.
21. McNulty H, McKinley MC, Wilson B, et al. Impaired functioning of thermolabile methylenetetrahydrofolate reductase is dependent on riboflavin status: implications for riboflavin requirements. *Am J Clin Nutr* 2002;76:436–41.
22. Stern LL, Mason JB, Selhub J, et al. Genomic DNA hypomethylation, a characteristic of most cancers, is present in peripheral leukocytes of individuals who are homozygous for the C677T polymorphism in the methylenetetrahydrofolate reductase gene. *Cancer Epidemiol Biomarkers Prev* 2000;9:849–53.
23. Narayanan S. The effect of folic acid and genetic polymorphisms on DNA stability and colorectal cancer. Aberdeen, Scotland: University of Aberdeen, 2001.
24. Friso S, Choi SW, Girelli D, et al. A common mutation in the 5,10-methylenetetrahydrofolate reductase gene affects genomic DNA methylation through an interaction with folate status. *Proc Natl Acad Sci U S A* 2002;99:5606–11.
25. Crott JW, Mashiyama ST, Ames BN, et al. Methylenetetrahydrofolate reductase C677T polymorphism does not alter folic acid deficiency-induced uracil incorporation into primary

TABLE 6. Research priorities

1. Further documentation of genotype frequencies: large, population-based studies of the polymorphisms reported in this paper and the additional, but less well studied, polymorphisms in these genes (e.g., *G1793A* in *MTHFR*), including prevalences of combinations of polymorphisms and prevalence in different age groups; particularly needed in non-White populations and less-investigated ethnic groups in the United States and Europe
2. Clarification of functional effects of the polymorphisms: including exploration of 1) consequences of carrying combinations of polymorphisms in both in vivo and in vitro systems and 2) in vivo functional effects of particular genotypes in persons with different levels of intake of folate and related dietary factors
3. Further investigation of hypothesized mechanisms: examination of whether the polymorphisms are associated, in humans, with genomic DNA methylation, uracil incorporation, or DNA strand breaks, including exploration of whether relations differ according to levels of folate and related dietary factors
4. Studies of gene-disease associations and gene-environment and gene-gene interactions: further large population-based studies of polymorphisms and cancer and adenomas, incorporating collection of high-quality dietary data and, ideally, blood biomarkers; these studies should be large enough to have sufficient power to investigate gene-environment and gene-gene interactions and to undertake subgroup analysis by age and ethnic group, of colon and rectal tumors, proximal and distal tumors, and tumors with microsatellite instability or loss of heterozygosity
5. Pooled analyses of studies of gene-disease associations and gene-environment and gene-gene interactions to facilitate subgroup analyses and investigation of interactions
6. Investigation of the role of other folate pathway genes, and interactions with alcohol-metabolizing genes, in the etiology of colorectal neoplasia
7. Investigation of genotype in adenomatous polyps: including 1) association with risk of recurrence and 2) association with particular pathologic features and 3) incorporation of genotyping in randomized controlled trials of folate supplementation in prevention of colorectal neoplasia
8. Further investigation of genotype and quality of life and the effectiveness of treatment in patients with colorectal cancers: large studies of representative groups of patients; analysis should include adjustment for known prognostic factors
9. Development of methodology for specifying hypotheses and statistical analysis in the context of interactions between multiple genes and multiple environmental factors

- human lymphocyte DNA in vitro. *Carcinogenesis* 2001;22:1019–25.
26. Andreassi MG, Botto N, Cocci F, et al. Methylenetetrahydrofolate reductase gene C677T polymorphism, homocysteine, vitamin B₁₂, and DNA damage in coronary artery disease. *Hum Genet* 2003;112:171–7.
 27. Zijno A, Andreoli C, Leopardi P, et al. Folate status, metabolic genotype, and biomarkers of genotoxicity in healthy subjects. *Carcinogenesis* 2003;24:1097–103.
 28. Chen J, Ma J, Stampfer MJ, et al. Linkage disequilibrium between the 677C>T and 1298A>C polymorphisms in human methylenetetrahydrofolate reductase gene and their contributions to risk of colorectal cancer. *Pharmacogenetics* 2002;12:339–42.
 29. Livers KJA, Boers GHJ, Verhoef P, et al. A second common variant in the methylenetetrahydrofolate reductase (*MTHFR*) gene and its relationship to *MTHFR* enzyme activity, homocysteine, and cardiovascular disease risk. *J Mol Med* 2001;79:522–8.
 30. Chango A, Boisson F, Barbé F, et al. The effect of 677C-T and 1298A-C mutations on plasma homocysteine and 5,10-methylenetetrahydrofolate reductase activity in healthy subjects. *Br J Nutr* 2000;83:593–6.
 31. Friedman G, Goldschmidt N, Friedlander Y, et al. A common mutation A1298C in human methylenetetrahydrofolate reductase gene: association with plasma total homocysteine and folate concentrations. *J Nutr* 1999;129:1656–61.
 32. van der Put NMJ, van der Molen EF, Kluijtmans LAJ, et al. Sequence analysis of the coding region of human methionine synthase: relevance to hyperhomocysteinemia in neural-tube defects and vascular disease. *QJM* 1997;90:511–17.
 33. Anwar W, Guéant JL, Abdelmoutaleb I, et al. Hyperhomocysteinemia is related to residual glomerular filtration and folate, but not to methylenetetrahydrofolate-reductase and methionine synthase polymorphisms, in supplemented end-stage renal disease patients undergoing hemodialysis. *Clin Chem Lab Med* 2001;39:747–52.
 34. Chen J, Stampfer MJ, Ma J, et al. Influence of a methionine synthase (D919G) polymorphism on plasma homocysteine and folate levels and relation to risk of myocardial infarction. *Atherosclerosis* 2001;154:667–72.
 35. Dekou V, Gudnason V, Hawe E, et al. Gene-environment and gene-gene interaction in the determination of plasma homocysteine levels in healthy middle-aged men. *Thromb Haemost* 2001;85:67–74.
 36. Silaste ML, Rantala M, Sampi M, et al. Polymorphisms of key enzymes in homocysteine metabolism affect diet responsiveness of plasma homocysteine in healthy women. *J Nutr* 2001;131:2643–7.
 37. Harmon DL, Shields DC, Woodside JV, et al. Methionine synthase D919G polymorphism is a significant but modest determinant of circulating homocysteine concentrations. *Genet Epidemiol* 1999;17:298–309.
 38. Gaughan DJ, Kluijtmans LAJ, Barbaux S, et al. The methionine synthase reductase (*MTRR*) A66G polymorphism is a novel genetic determinant of plasma homocysteine concentrations. *Atherosclerosis* 2001;157:451–6.
 39. Geisel J, Zimbelmann I, Schorr H, et al. Genetic defects as important factors for moderate hyperhomocysteinemia. *Clin Chem Lab Med* 2001;39:698–704.
 40. O'Leary VB, Parle-McDermott A, Molloy AM, et al. *MTRR* and *MTHFR* polymorphism: link to Down syndrome? *Am J Med Genet* 2002;107:151–5.
 41. Kraus JP. Biochemistry and molecular genetics of cystathionine β -synthase deficiency. *Eur J Pediatr* 1998;157(suppl 2):S50–S53.
 42. Kluijtmans LAJ, Boers GHJ, Trijbels FJM, et al. A common 844INS68 insertion variant in the cystathionine β -synthase gene. *Biochem Mol Med* 1997;62:23–5.
 43. Tsai MY, Yang F, Bignell M, et al. Relation between plasma homocysteine concentration, the 844ins68 variant of the cys-

- tathionine β -synthase gene, and pyridoxal-5'-phosphate concentration. *Mol Genet Metab* 1999;67:352-6.
44. Luo HR, Lü XM, Yao YG, et al. Length polymorphism of thymidylate synthase regulatory region in Chinese populations and evolution of the novel alleles. *Biochem Genet* 2002;40:41-51.
 45. Marsh S, Ameyaw MM, Githang'a J, et al. Novel thymidylate synthase enhancer region alleles in African populations. *Hum Mutat* 2000 Dec;16:528.
 46. Horie N, Aiba H, Oguro K, et al. Functional analysis and DNA polymorphism of the tandemly repeated sequences in the 5'-terminal regulatory region of the human gene for thymidylate synthase. *Cell Struct Funct* 1995;20:191-7.
 47. Trinh BN, Ong CN, Coetzee GA, et al. Thymidylate synthase: a novel genetic determinant of plasma homocysteine and folate levels. *Hum Genet* 2002;111:299-302.
 48. Ulrich CM, Bigler J, Velicer CM, et al. Searching expressed sequence tag databases: discovery and confirmation of a common polymorphism in the thymidylate synthase gene. *Cancer Epidemiol Biomarkers Prev* 2000;9:1381-5.
 49. Botto LD, Yang Q. 5,10-Methylenetetrahydrofolate reductase gene variants and congenital anomalies: a HuGE review. *Am J Epidemiol* 2000;151:862-77.
 50. Robien K, Ulrich CM. 5,10-Methylenetetrahydrofolate reductase polymorphisms and leukemia risk: a HuGE minireview. *Am J Epidemiol* 2003;157:571-82.
 51. Barber R, Shalat S, Hendricks K, et al. Investigation of folate pathway gene polymorphisms and the incidence of neural tube defects in a Texas Hispanic population. *Mol Genet Metab* 2000;70:45-52.
 52. Volcik KA, Blanton SH, Tyerman GH, et al. Methylenetetrahydrofolate reductase and spina bifida: evaluation of level of defect and maternal genotypic risk in Hispanics. *Am J Med Genet* 2000;95:21-7.
 53. Peng F, Labelle LA, Rainey BJ, et al. Single nucleotide polymorphisms in the methylenetetrahydrofolate reductase gene are common in US Caucasian and Hispanic American populations. *Int J Mol Med* 2001;8:509-11.
 54. Le Marchand L, Donlon T, Hankin JH, et al. B-vitamin intake, metabolic genes, and colorectal cancer risk (United States). *Cancer Causes Control* 2002;13:239-48.
 55. Sharp L, Little J, Schofield AC, et al. Folate and breast cancer: the role of polymorphisms in methylenetetrahydrofolate reductase (MTHFR). *Cancer Lett* 2002;181:65-71.
 56. Sharp L, Little J, Brockton N, et al. Dietary intake of folate and related micronutrients, genetic polymorphisms in MTHFR and colorectal cancer: a population-based case-control study in Scotland. (Abstract). *J Nutr* 2002;132(11S):3542S.
 57. Brunelli das Neves Grillo L, Acácio GL, Barini R, et al. Mutations in the methylene-tetrahydrofolate reductase gene and Down syndrome. *Cad Saude Publica* 2002;18:1795-7.
 58. Akar N, Akar E, Akcay R, et al. Effect of methylenetetrahydrofolate reductase 677 C-T, 1298 A-C, and 1317 T-C on factor V 1691 mutation in Turkish deep vein thrombosis patients. *Thromb Res* 2000;91:163-7.
 59. Gebhardt GS, Scholtz CL, Hillermann R, et al. Combined heterozygosity for methylenetetrahydrofolate reductase (MTHFR) mutations C677T and A1298C is associated with abruptio placentae but not with intrauterine growth restriction. *Eur J Obstet Gynecol Reprod Biol* 2001;97:174-7.
 60. Stegmann K, Ziegler A, Ngo ETKM, et al. Linkage disequilibrium of MTHFR genotypes 677C/T-1298A/C in the German population and association studies in probands with neural tube defects (NTD). *Am J Med Genet* 1999;87:23-9.
 61. Shen H, Spitz MR, Wang LE, et al. Polymorphisms of methylene-tetrahydrofolate reductase and risk of lung cancer: a case-control study. *Cancer Epidemiol Biomarkers Prev* 2001;10:397-401.
 62. Meisel C, Cascorbi I, Gerloff T, et al. Identification of six methylenetetrahydrofolate reductase (MTHFR) genotypes resulting from common polymorphisms: impact on plasma homocysteine levels and development of coronary artery disease. *Atherosclerosis* 2001;154:651-8.
 63. Skibola CF, Smith MT, Kane E, et al. Polymorphisms in the methylenetetrahydrofolate reductase gene are associated with susceptibility to acute leukemia in adults. *Proc Natl Acad Sci U S A* 1999;96:12810-15.
 64. Song C, Xing D, Tan W, et al. Methylenetetrahydrofolate reductase polymorphisms increase risk of esophageal squamous cell carcinoma in a Chinese population. *Cancer Res* 2001;61:3272-5.
 65. Zhang G, Dai C. Gene polymorphisms of homocysteine metabolism-related enzymes in Chinese patients with occlusive coronary artery or cerebral vascular diseases. *Thromb Res* 2001;104:187-95.
 66. Morita S, Yano M, Tsujinaka T, et al. Genetic polymorphisms of drug-metabolizing enzymes and susceptibility to head-and-neck squamous-cell carcinoma. *Int J Cancer* 1999;80:685-8.
 67. Matsuo K, Suzuki R, Hamajima N, et al. Association between polymorphisms of folate- and methionine-metabolizing enzymes and susceptibility to malignant lymphoma. *Blood* 2001;97:3205-9.
 68. Jang Y, Park HY, Lee JH, et al. A polymorphism of the methylenetetrahydrofolate reductase and methionine synthase gene in CAD patients: association with plasma folate, vitamin B₁₂ and homocysteine. *Nutr Res* 2002;22:965-76.
 69. Feix A, Fritsche-Polanz R, Kletzmayer J, et al. Increased prevalence of combined MTR and MTHFR genotypes among individuals with severely elevated total homocysteine plasma levels. *Am J Kidney Dis* 2001;38:956-64.
 70. Kimura F, Franke KH, Steinhoff C, et al. Methyl group metabolism gene polymorphisms and susceptibility to prostatic carcinoma. *Prostate* 2000;45:225-31.
 71. Kimura F, Florl AR, Steinhoff C, et al. Polymorphic methyl group metabolism genes in patients with transitional cell carcinoma of the urinary bladder. *Mutat Res* 2001;458:49-54.
 72. D'Angelo A, Coppola A, Madonna P, et al. The role of vitamin B₁₂ in fasting hyperhomocysteinemia and its interaction with the homozygous C677T mutation of the methylenetetrahydrofolate reductase (MTHFR) gene. *Thromb Haemost* 2000;83:563-70.
 73. De Marco P, Calevo MG, Moroni A, et al. Study of MTHFR and MS polymorphisms as risk factors for NTD in the Italian population. *J Hum Genet* 2002;47:319-24.
 74. Lucock M, Daskalakis I, Briggs D, et al. Altered folate metabolism and disposition in mothers affected by a spina bifida pregnancy: influence of 677C→T methylenetetrahydrofolate reductase and 2756A→G methionine synthase genotypes. *Mol Genet Metab* 2000;70:27-44.
 75. Christensen B, Arbour L, Tran P, et al. Genetic polymorphisms in methylenetetrahydrofolate reductase and methionine synthase, folate levels in red blood cells, and risk of neural tube defects. *Am J Med Genet* 1999;84:151-7.
 76. Chen J, Giovannucci E, Hankinson SE, et al. A prospective study of methylenetetrahydrofolate reductase and methionine synthase gene polymorphisms, and risk of colorectal adenoma. *Carcinogenesis* 1998;19:2129-32.
 77. Barbaux S, Plomin R, Whitehead AS. Polymorphisms of genes controlling homocysteine/folate metabolism and cognitive function. *Neuroreport* 2000;11:1133-6.

78. Tsai MY, Welge BG, Hanson NQ, et al. Genetic causes of mild hyperhomocysteinemia in patients with premature occlusive coronary artery diseases. *Atherosclerosis* 1999;143:163–70.
79. Tsai MY, Bignell M, Yang F, et al. Polygenic influence on plasma homocysteine: association of two prevalent mutations, the 844ins68 of cystathionine β -synthase and A2756G of methionine synthase, with lowered plasma homocysteine levels. *Atherosclerosis* 2000;149:131–7.
80. Johanning GL, Tamura T, Johnston KE, et al. Comorbidity of 5,10-methylenetetrahydrofolate reductase and methionine synthase gene polymorphisms and risk for neural tube defects. *J Med Genet* 2000;37:949–51.
81. Conroy JM, Trivedi G, Sovd T, et al. The allele frequency of mutations in four genes that confer enhanced susceptibility to venous thromboembolism in an unselected group of New York State newborns. *Thromb Res* 2000;99:317–24.
82. Wang XL, Duarte N, Cai H, et al. Relationship between total plasma homocysteine, polymorphisms of homocysteine metabolism related enzymes, risk factors and coronary artery disease in the Australian hospital-based population. *Atherosclerosis* 1999;146:133–40.
83. Ray JG, Langman LJ, Vermeulen MJ, et al. Genetics University of Toronto Thrombophilia Study in Women (GUTTSI): genetic and other risk factors for venous thromboembolism in women. *Curr Control Trials Cardiovasc Med* 2001;2:141–9.
84. Hobbs CA, Sherman SL, Yi P, et al. Polymorphisms in genes involved in folate metabolism as maternal risk factors for Down syndrome. *Am J Hum Genet* 2000;67:623–30.
85. Hassold TJ, Burrage LC, Chan ER, et al. Maternal folate polymorphisms and the etiology of human nondisjunction. *Am J Hum Genet* 2001;69:434–9.
86. Franco RF, Elion J, Lavinha J, et al. Heterogeneous ethnic distribution of the 844ins68 in the cystathionine β -synthase gene. *Hum Hered* 1998;48:338–42.
87. Orendác M, Musková B, Richterová E, et al. Is the common 844ins68 polymorphism in the cystathionine β -synthase gene associated with atherosclerosis? *J Inherit Metab Dis* 1999;22:674–5.
88. Richter B, Stegmann K, Roper B, et al. Interaction of folate and homocysteine pathway genotypes evaluated in susceptibility to neural tube defects (NTD) in a German population. *J Hum Genet* 2001;46:105–9.
89. Linnebank M, Homberger A, Junker R, et al. High prevalence of the 1278T mutation of the human cystathionine β -synthase detected by a novel screening application. *Thromb Haemost* 2001;85:986–8.
90. Grossmann R, Geisen U, Merati G, et al. Genetic risk factors in young adults with 'cryptogenic' ischemic cerebrovascular disease. *Blood Coagul Fibrinolysis* 2002;13:583–90.
91. Sperandio MP, de Franchis R, Andria G, et al. A 68-bp insertion found in a homocystinuric patient is a common variant and is skipped by alternative splicing of the cystathionine β -synthase mRNA. *Am J Hum Genet* 1996;59:1393–5.
92. Giusti B, Camacho-Vanegas O, Attanasio M, et al. Microheterogeneity in the distribution of the 844ins68 in the cystathionine β -synthase gene in Italy. *Thromb Res* 1999;94:249–54.
93. Papa A, De Stefano V, Danese S, et al. Hyperhomocysteinemia and prevalence of polymorphisms of homocysteine metabolism-related enzymes in patients with inflammatory bowel disease. *Am J Gastroenterol* 2001;96:2677–82.
94. de Franchis R, Fermo I, Mazzola G, et al. Contribution of the cystathionine β -synthase gene (844ins68) polymorphism to the risk of early-onset venous and arterial occlusive disease and of fasting hyperhomocysteinemia. *Thromb Haemost* 2000;84:576–82.
95. Olivieri O, Friso S, Trabetti E, et al. Homocysteine and atherosclerotic renal artery stenosis. *Clin Exp Med* 2001;1:211–18.
96. Folsom AR, Nieto FJ, McGovern PG, et al. Prospective study of coronary heart disease incidence in relation to fasting total homocysteine, related genetic polymorphisms, and B vitamins. *Circulation* 1998;98:204–10.
97. Tsai MY, Bignell M, Schwichtenberg K, et al. High prevalence of a mutation in the cystathionine β -synthase gene. *Am J Hum Genet* 1996;59:1262–7.
98. Shannon B, Gnanasampathan S, Beilby J, et al. A polymorphism in the methylenetetrahydrofolate reductase gene predisposes to colorectal cancers with microsatellite instability. *Gut* 2002;50:520–4.
99. Chen J, Hunter DJ, Stampfer MJ, et al. A novel polymorphism in the thymidylate synthase gene promoter influences plasma folate level and may modify the risk of colorectal cancer in a prospective study. (Abstract). Presented at the 93rd Annual Meeting of the American Association for Cancer Research, San Francisco, California, April 6–10, 2002.
100. Marsh S, Collie-Duguid ESR, Li T, et al. Ethnic variation in the thymidylate synthase enhancer region polymorphism among Caucasian and Asian populations. *Genomics* 1999;58:310–12.
101. Skibola CF, Smith MT, Hubbard A, et al. Polymorphisms in the thymidylate synthase and serine hydroxymethyltransferase genes and risk of adult acute lymphocytic leukemia. *Blood* 2002;99:3786–91.
102. Ulrich CM, Bigler J, Bostick R, et al. Thymidylate synthase promoter polymorphism, interaction with folate intake, and risk of colorectal adenomas. *Cancer Res* 2002;62:3361–4.
103. de Franchis R, Botto LD, Sebastio G, et al. Spina bifida and folate-related genes: a study of gene-gene interactions. *Genet Med* 2002;4:126–30.
104. Xu J, Turner A, Little J, et al. Positive results in association studies are associated with departure from Hardy-Weinberg equilibrium: hint for genotyping error? *Hum Genet* 2002;111:573–4.
105. Garte S. The role of ethnicity in cancer susceptibility gene polymorphisms: the example of CYP1A1. *Carcinogenesis* 1998;19:1329–32.
106. Ferlay J, Bray F, Pisani P, et al. GLOBOCAN 2000: cancer incidence, mortality and prevalence worldwide, version 1.0. IARC CancerBase no. 5. Lyon, France: International Agency for Research on Cancer, 2001.
107. Parkin DM, Pisani P, Ferlay J. Estimates of the worldwide incidence of 25 major cancers in 1990. *Int J Cancer* 1999;80:827–41.
108. Parkin DM, Whelan SL, Ferlay J, et al. Cancer incidence in five continents. Vol VII. Lyon, France: International Agency for Research on Cancer, 1997. (IARC scientific publication no. 143).
109. Cotton S, Sharp L, Little J. The adenoma-carcinoma sequence and prospects for the prevention of colorectal neoplasia. *Crit Rev Oncog* 1996;7:293–342.
110. Hawkins NJ, Ward RL. Sporadic colorectal cancers with microsatellite instability and their possible origin in hyperplastic polyps and serrated adenomas. *J Natl Cancer Inst* 2001;93:1307–13.
111. Mecklin JP, Ponz de Leon M. Epidemiology of HNPCC. *Anticancer Res* 1994;14:1625–9.
112. St John DJB, McDermott FT, Hopper JL, et al. Cancer risk in relatives of patients with common colorectal cancer. *Ann Intern Med* 1993;118:785–90.
113. Fuchs CS, Giovannucci EL, Colditz GA, et al. A prospective study of family history and the risk of colorectal cancer. *N Engl J Med* 1994;331:1669–74.

114. Winawer SJ, Zauber AG, Gerdes H, et al. Risk of colorectal cancer in the families of patients with adenomatous polyps. *N Engl J Med* 1996;334:82–7.
115. Khoury MJ, Beaty TH, Liang KY. Can familial aggregation of disease be explained by familial aggregation of environmental risk factors? *Am J Epidemiol* 1988;127:674–83.
116. Haenszel W, Kurihara M. Studies of Japanese migrants: mortality from cancer and other diseases among Japanese in the United States. *J Natl Cancer Inst* 1968;40:43–68.
117. Bergström A, Pisani V, Tenet V, et al. Overweight as an avoidable cause of cancer in Europe. *Int J Cancer* 2001;91:421–30.
118. The role of weight control and physical activity in cancer prevention. IARC Working Group on the Evaluation of Cancer. IARC handbooks of cancer prevention. Vol 6. Lyon, France: International Agency for Research on Cancer, 2002.
119. Giovannucci E. An updated review of the epidemiological evidence that cigarette smoking increases risk of colorectal cancer. *Cancer Epidemiol Biomarkers Prev* 2001;10:725–31.
120. World Cancer Research Fund/American Institute for Cancer Research. Food, nutrition and the prevention of cancer: a global perspective. Menasha, WI: Banta Book Group, 1997.
121. Beral V, Banks E, Reeves G, et al. Use of HRT and the subsequent risk of cancer. *J Epidemiol Biostat* 1999;4:191–215.
122. Rossouw JE, Anderson GL, Prentice RL, et al. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results from the Women's Health Initiative randomized controlled trial. *JAMA* 2002;288:321–33.
123. Non-steroidal anti-inflammatory drugs. IARC Working Group on the Evaluation of Cancer Preventive Agents. IARC handbooks of cancer prevention. Vol 1. Lyon, France: International Agency for Research on Cancer, 1997.
124. Terry P, Giovannucci E, Michels KB, et al. Fruit, vegetables, dietary fiber and risk of colorectal cancer. *J Natl Cancer Inst* 2001;93:525–33.
125. Flood A, Velie EM, Chatterjee N, et al. Fruit and vegetable intakes and the risk of colorectal cancer in the Breast Cancer Detection Demonstration Project follow-up cohort. *Am J Clin Nutr* 2002;75:936–43.
126. Glynn SA, Albanes D, Pietinen P, et al. Colorectal cancer and folate status: a nested case-control study among male smokers. *Cancer Epidemiol Biomarkers Prev* 1996;5:487–94.
127. Kato I, Dnistrian AM, Schwartz M, et al. Serum folate, homocysteine and colorectal cancer risk in women: a nested case-control study. *Br J Cancer* 1999;79:1917–21.
128. Bird CL, Swendseid ME, Witte JS, et al. Red cell and plasma folate, folate consumption, and the risk of colorectal adenomatous polyps. *Cancer Epidemiol Biomarkers Prev* 1995;4:709–14.
129. Paspatis GA, Kalafatis E, Oros L, et al. Folate status and adenomatous colonic polyps: a colonoscopically controlled study. *Dis Colon Rectum* 1995;38:64–8.
130. Giovannucci E, Rimm EB, Ascherio A, et al. Alcohol, low-methionine–low-folate diets, and risk of colon cancer in men. *J Natl Cancer Inst* 1995;87:265–73.
131. Giovannucci E, Stampfer MJ, Colditz GA, et al. Multivitamin use, folate, and colon cancer in women in the Nurses' Health Study. *Ann Intern Med* 1998;129:517–24.
132. Su LJ, Arab L. Nutritional status of folate and colon cancer risk: evidence from NHANES I epidemiologic follow-up study. *Ann Epidemiol* 2001;11:65–72.
133. Terry P, Jain M, Miller AB, et al. Dietary intake of folic acid and colorectal cancer risk in a cohort of women. *Int J Cancer* 2002;97:864–7.
134. Freudenheim JL, Graham S, Marshall JR, et al. Folate intake and carcinogenesis of the colon and rectum. *Int J Epidemiol* 1991;20:368–74.
135. Benito E, Stiggelbout A, Bosch FX, et al. Nutritional factors in colorectal cancer risk: a case-control study in Majorca. *Int J Cancer* 1991;49:161–7.
136. Meyer F, White E. Alcohol and nutrients in relation to colon cancer in middle-aged adults. *Am J Epidemiol* 1993;138:225–36.
137. Ferraroni M, La Vecchia C, D'Avanzo B, et al. Selected micronutrient intake and the risk of colorectal cancer. *Br J Cancer* 1994;70:1150–5.
138. Boutron-Ruault MC, Senesse P, Faivre J, et al. Folate and alcohol intakes: related or independent roles in the adenoma-carcinoma sequence? *Nutr Cancer* 1996;26:337–46.
139. La Vecchia C, Braga C, Negri E, et al. Intake of selected micronutrients and risk of colorectal cancer. *Int J Cancer* 1997;73:525–30.
140. Slattery ML, Schaffer D, Edwards SL, et al. Are dietary factors involved in DNA methylation associated with colon cancer? *Nutr Cancer* 1997;28:52–62.
141. Levi F, Pasche C, Lucchini F, et al. Selected micronutrients and colorectal cancer: a case-control study from the Canton of Vaud, Switzerland. *Eur J Cancer* 2000;36:2115–19.
142. Benito E, Cabeza E, Moreno V, et al. Diet and colorectal adenomas: a case-control study in Majorca. *Int J Cancer* 1993;55:213–19.
143. Giovannucci E, Stampfer MJ, Colditz GA, et al. Folate, methionine and alcohol intake and risk of colorectal adenoma. *J Natl Cancer Inst* 1993;85:875–84.
144. Tseng M, Murray SC, Kupper LL, et al. Micronutrients and the risk of colorectal adenomas. *Am J Epidemiol* 1996;144:1005–14.
145. Breuer-Katschinski B, Nemes K, Marr A, et al. Colorectal adenomas and diet: a case-control study. *Dig Dis Sci* 2001;46:86–95.
146. Baron JA, Sandler RS, Haile RW, et al. Folate intake, alcohol consumption, cigarette smoking, and risk of colorectal adenomas. *J Natl Cancer Inst* 1998;90:57–62.
147. Paspatis G, Xourgias B, Mylonakou E, et al. A prospective clinical trial to determine the influence of folate supplementation on the formation of recurrent colonic adenomas. (Abstract). *Gastroenterology* 1994;106:A425.
148. Herbert V. Recommended dietary intakes (RDI) of folate in humans. *Am J Clin Nutr* 1987;45:661–70.
149. Chen J, Giovannucci E, Kelsy K, et al. A methylenetetrahydrofolate reductase polymorphism and the risk of colorectal cancer. *Cancer Res* 1996;56:4862–4.
150. Slattery ML, Potter JD, Samowitz W, et al. Methylenetetrahydrofolate reductase, diet, and risk of colon cancer. *Cancer Epidemiol Biomarkers Prev* 1999;8:513–18.
151. Park KS, Mok JW, Kim JC. The 677>T mutation in 5,10-methylenetetrahydrofolate reductase and colorectal cancer risk. *Genet Test* 1999;3:233–6.
152. Delgado-Enciso I, Martinez-Garza SG, Rojas-Martinez A, et al. 677T mutation of the MTHFR gene in adenomas and colorectal cancer in a population sample from the northeastern Mexico. *Rev Gastroenterol Mex* 2001;66:32–7.
153. Sharp L, Little J, Brockton N, et al. Genetic polymorphisms in folate metabolism, dietary folate intake and colorectal cancer: a population-based case-control study. (Abstract). *J Epidemiol Community Health* 2001;55:A27.
154. Keku T, Millikan R, Worley K, et al. 5,10-methylenetetrahydrofolate reductase codon 677 and 1298 polymorphisms and colon cancer in African Americans and whites. *Cancer Epidemiol Biomarkers Prev* 2002;11:1611–21.
155. Sachse C, Smith G, Wilkie MJV, et al. A pharmacogenetic study to investigate the role of dietary carcinogens in the etiology of colorectal cancer. *Carcinogenesis* 2002;23:1839–49.

156. Ioannidis JP, Ntzani EE, Trikalinos TA, et al. Replication validity of genetic association studies. *Nat Genet* 2001;29:306–9.
157. Lohmueller KE, Pearce CL, Pike M, et al. Meta-analysis of genetic association studies supports a contribution of common variants to susceptibility to common disease. *Nat Genet* 2003;33:177–82.
158. Klerk M, Verhoef P, Clarke R, et al. MTHFR 677C→T polymorphism and risk of coronary heart disease. *JAMA* 2002;288:2023–31.
159. Ulrich CM, Kampman E, Bigler J, et al. Colorectal adenomas and the C677T MTHFR polymorphism: evidence for gene-environment interaction? *Cancer Epidemiol Biomarkers Prev* 1999;8:659–68.
160. Levine AJ, Siegmund KD, Ervin CM, et al. The methylenetetrahydrofolate reductase 677C→T polymorphism and distal colorectal adenoma risk. *Cancer Epidemiol Biomarkers Prev* 2000;9:657–63.
161. Marugame T, Tsuji E, Inoue H, et al. Methylenetetrahydrofolate reductase polymorphism and risk of colorectal adenomas. *Cancer Lett* 2000;151:181–6.
162. Ulvik A, Evensen ET, Lien EA, et al. Smoking, folate and methylenetetrahydrofolate reductase status as interactive determinants of adenomatous and hyperplastic polyps of colorectum. *Am J Med Genet* 2001;101:246–54.
163. Ulrich CM, Kampman E, Bigler J, et al. Lack of association between the C677T MTHFR polymorphism and colorectal hyperplastic polyps. *Cancer Epidemiol Biomarkers Prev* 2000;9:427–34.
164. Lenz HJ, Zhang W, Zahedy S, et al. A 6 base-pair deletion in the 3' UTR of the thymidylate synthase (TS) gene predicts TS mRNA expression in colorectal tumours: a possible candidate gene for colorectal cancer risk. (Abstract). Presented at the 93rd Annual Meeting of the American Association for Cancer Research, San Francisco, California, April 6–10, 2002.
165. Ramsbottom D, Scott JM, Molloy A, et al. Are common mutations of cystathionine β -synthase involved in the aetiology of neural tube defects? *Clin Genet* 1997;51:39–42.
166. Mills JL, Kirke PN, Molloy AM, et al. Methylenetetrahydrofolate reductase thermolabile variant and oral clefts. *Am J Med Genet* 1999;86:71–4.
167. Shen H, Xu Y, Zheng Y, et al. Polymorphisms of 5,10 methylenetetrahydrofolate reductase and risk of gastric cancer in a Chinese population: a case-control study. *Int J Cancer* 2001;95:332–6.
168. Dilley A, Hooper WC, El-Jamil M, et al. Mutations in the genes regulating methylene tetrahydrofolate reductase (MTHFR C-T677) and cystathionine β -synthase (CBS G-A919, CBS T-c833) are not associated with myocardial infarction in African Americans. *Thromb Res* 2001;103:109–15.
169. Verhaar MC, Stroes E, Rabelink TJ. Folate and cardiovascular disease. *Arterioscler Thromb Vasc Biol* 2002;22:6–13.
170. Molloy AM, Scott JM. Folate and prevention of disease. *Public Health Nutr* 2001;4:601–9.
171. Hustad S, Ueland PM, Vollset SE, et al. Riboflavin as a determinant of plasma total homocysteine: effect modification by the methylenetetrahydrofolate reductase C677T polymorphism. *Clin Chem* 2000;46:1065–71.
172. Wolf CR, Smith G. Cytochrome P450 CYP2D6. *IARC Sci Publ* 1999;209–29.
173. Little J, Bradley L, Bray MS, et al. Reporting, appraising, and integrating data on genotype prevalence and gene-disease associations. *Am J Epidemiol* 2002;156:300–10.
174. Genelex Corporation. Health and DNA. Nutritional genetics (Richmond, Washington), 1995–2002. (<http://www.healthanddna.com/nutrigeneticstest.html>).
175. Sciona Ltd. Body benefits—nutrition. Havant, United Kingdom, 2003 (<http://www.sciona.com/coresite/products/nutrition7.htm>).
176. Wisotzkey JD, Toman J, Bell T, et al. MTHFR (C677T) polymorphisms and stage III colon cancer: response to therapy. *Mol Diagn* 1999;4:95–9.
177. Edler D, Hallström M, Johnston PG, et al. Thymidylate synthase expression: an independent prognostic factor for local recurrence, distant metastasis, disease-free and overall survival in rectal cancer. *Clin Cancer Res* 2000;6:1378–84.
178. Aschele C, Debernardis D, Bandelloni R, et al. Thymidylate synthase protein expression in colorectal cancer metastases predicts for clinical outcome to leucovorin-modulated bolus or infusional 5-fluorouracil but not methotrexate-modulated bolus 5-fluorouracil. *Ann Oncol* 2002;13:1882–92.
179. Etienne MC, Chazal M, Laurent-Puig P, et al. Prognostic value of tumoral thymidylate synthase and p53 in metastatic colorectal cancer patients receiving fluorouracil-based chemotherapy: phenotypic and genotypic analyses. *J Clin Oncol* 2002;20:2832–43.
180. Allegra CJ, Parr AL, Wold LE, et al. Investigation of the prognostic and predictive value of thymidylate synthase, p53, and Ki-67 in patients with locally advanced colon cancer. *J Clin Oncol* 2002;20:1735–43.
181. Sugiyama Y, Kato T, Nakazato H, et al. Retrospective study on thymidylate synthase as a predictor of outcome and sensitivity to adjuvant chemotherapy in patients with curatively resected colorectal cancer. *Anticancer Drugs* 2002;13:931–8.
182. Iacopetta B, Grieco F, Joseph D, et al. A polymorphism in the enhancer region of the thymidylate synthase promoter influences the survival of colorectal cancer patients treated with 5-fluorouracil. *Br J Cancer* 2001;85:827–30.
183. Marsh S, McKay JA, Cassidy J, et al. Polymorphism in the thymidylate synthase promoter enhancer region in colorectal cancer. *Int J Oncol* 2001;19:383–6.
184. Pullarkat ST, Stoehlmacher J, Ghaderi V, et al. Thymidylate synthase gene polymorphism determines response and toxicity of 5-FU chemotherapy. *Pharmacogenomics J* 2001;1:65–70.
185. Villafranca E, Okruzhnov Y, Dominguez MA, et al. Polymorphisms of the repeated sequences in the enhancer region of the thymidylate synthase gene promoter may predict downstaging after preoperative chemoradiation in rectal cancer. *J Clin Oncol* 2001;19:1779–86.
186. Ulrich CM, Robien K, Sparks R. Pharmacogenetics and folate metabolism—a promising direction. *Pharmacogenetics* 2002;3:299–313.

APPENDIX. Internet sites pertaining to folate metabolism and colorectal neoplasia**Data on cancer incidence, survival, and mortality**

International Agency for Research on Cancer (IARC)—Cancer Mondial:

<http://www-dep.iarc.fr/dataava/infodata.htm>

Surveillance, Epidemiology, and End Results (SEER) Program:

<http://www.seer.cancer.gov/publicdata/>

National Program of Cancer Registries (NPCR):

<http://www.cdc.gov/cancer/npcr>

Information on cancer

Cancer Research UK:

<http://www.cancerresearchuk.org/>

National Cancer Institute—cancer.gov:

<http://www.nci.nih.gov/>

American Cancer Society:

<http://www.cancer.org/docroot/home/index.asp>

Genetics information

Human Genome Epidemiology Network (HuGENet):

<http://www.cdc.gov/genomics/hugenet/default.htm>

Public Health Genetics Unit:

<http://www.medschl.cam.ac.uk/phgu/>

Online Mendelian Inheritance in Man (OMIM):

<http://www3.ncbi.nlm.nih.gov/Omim/searchomim.html>

GenAtlas:

<http://www.dsi.univ-paris5.fr/genatlas>

GeneCards:

<http://www.cgal.icnet.uk/genecards>

National Center for Biotechnology Information:

<http://www.ncbi.nlm.nih.gov/>

UK Human Genome Mapping Project (includes links to other sites via The Genome Web):

<http://www.hgmp.mrc.ac.uk/>